



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2021

The Scontrone turtles – A new insular testudinoid fauna from the late Miocene of the Central Mediterranean

Georgalis, Georgios L ; Delfino, Massimo

Abstract: We here describe a small turtle assemblage originating from the early Tortonian (late Miocene) palaeoisland of Scontrone, central Italy, a locality previously known mostly for its endemic mammals and giantbirds, which were otherwise shared only with the Gargano localities, another fossiliferous area belonging to the same palaeobioprovince. The fossil turtle remains from Scontrone are referred to the geoemydid *Mauremys* sp. and a so far unidentified large-sized testudinid. The biogeographic origins of the Scontrone insular chelonians are discussed. The Scontrone geoemydid adds to the known occurrences of *Mauremys* in the late Miocene of the Mediterranean. The Scontrone large tortoise represents the oldest known Mediterranean insular testudinid, predating significantly the well-known Quaternary endemic island tortoises of the area.

DOI: <https://doi.org/10.1016/j.geobios.2021.05.001>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-204606>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.

Originally published at:

Georgalis, Georgios L; Delfino, Massimo (2021). The Scontrone turtles – A new insular testudinoid fauna from the late Miocene of the Central Mediterranean. *Geobios*, 68:71-81.

DOI: <https://doi.org/10.1016/j.geobios.2021.05.001>



Original article

The Scontrone turtles – A new insular testudinoid fauna from the late Miocene of the Central Mediterranean ☆

Georgios L. Georgalis ^{a,*}, Massimo Delfino ^{b,c,*}^a Palaeontological Institute and Museum, University of Zurich, Karl Schmid-Strasse 4, 8006 Zurich, Switzerland^b Dipartimento di Scienze della Terra, Università di Torino, 35, via Valperga Caluso, 10125 Torino, Italy^c Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona, Edifici ICTA -ICP, Carrer de les Columnes s/n, Campus de la UAB, 08193 Cerdanyola del Vallès, Barcelona, Spain

ARTICLE INFO

Article history:

Received 2 January 2021

Accepted 15 May 2021

Available online 10 June 2021

Keywords:

Testudines
Geoemydidae
Testudinidae
Palaeoisland
Biogeography
Miocene

ABSTRACT

We here describe a small turtle assemblage originating from the early Tortonian (late Miocene) palaeoisland of Scontrone, central Italy, a locality previously known mostly for its endemic mammals and giant birds, which were otherwise shared only with the Gargano localities, another fossiliferous area belonging to the same palaeobioprovince. The fossil turtle remains from Scontrone are referred to the geoemydid *Mauremys* sp. and a so far unidentified large-sized testudinid. The biogeographic origins of the Scontrone insular chelonians are discussed. The Scontrone geoemydid adds to the known occurrences of *Mauremys* in the late Miocene of the Mediterranean. The Scontrone large tortoise represents the oldest known Mediterranean insular testudinid, predating significantly the well-known Quaternary endemic island tortoises of the area.

© 2021 The Author(s). Published by Elsevier Masson SAS. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Mediterranean islands offer a unique insight into the evolution of several vertebrate lineages and witness the acquisition of extreme and astonishing insular adaptations and species radiations (Van der Geer et al., 2010; Herridge and Lister, 2012; Palombo et al., 2013; Georgalis et al., 2019). Especially for testudinid turtles, Mediterranean islands are long known to have drastically affected the evolution of this lineage, witnessing the radiation and extinction of an array of unique, endemic, and bizarre forms across different insular ecosystems (Leith Adams, 1877; Bate, 1914; Abbazzi et al., 2008; Luján et al., 2017).

Besides the islands that are present today in the Mediterranean Sea, other palaeoislands also existed in the area during its Cenozoic past, that are currently, however, part of mainland Europe (Lanza, 1984). The most prominent and well known among such cases represent the famous fossiliferous localities in Gargano in southern Italy and Scontrone in central Italy, which were during the late Miocene, parts of the Abruzzo-Apulia Platform that has been repeatedly isolated from neighbouring mainland areas for long

periods of time (Patacca et al., 2008a, 2008b, 2013). These two localities have been grouped in the same palaeobioprovince, the Apulo-Abruzzi area (Rook et al., 2008). The majority of the different fissures of Gargano pertain to a timespan between the late Miocene and the Pliocene and have yielded a considerably rich fossil fauna over the past decades (Freudenthal, 1971, 1972, 1976, 1985; Ballmann, 1976; Leinders, 1984; Abbazzi et al., 1996; Delfino et al., 2007; Masini et al., 2010, 2013; Masini and Fanfani, 2013; Freudenthal et al., 2013; Villier and Carnevale, 2013; Meijer, 2014; Van Der Geer, 2014; Savorelli and Masini, 2016; Pavia et al., 2017; Pavia, 2013); younger, Early Pleistocene sites, known as Pirro Nord are also known in Gargano (for the reptiles see, among others, Delfino and Bailon, 2000; Blain et al., 2019). Scontrone is older (Tortonian) and has yielded a poorer, though still unique faunal assemblage, comprising also endemic mammals and birds with peculiar morphologies (Rustioni et al., 1992; Mazza and Rustioni, 1996, 2008, 2011; Savorelli et al., 2017; Meijer, 2014; Van Der Geer, 2014; Pavia et al., 2017).

Turtle remains from these unique insular environments of the late Neogene palaeoislands of Gargano and Scontrone have been only poorly known, being solely represented by preliminary descriptions in a thesis of scarce testudinid remains from Gargano (Delfino, 2002) and simple mentions (Mazza and Rustioni, 2008; Patacca et al., 2013; Collareta et al., 2020) or partial descriptions in a thesis (Chesi, 2009b) of *Mauremys* from Scontrone. Here we

☆ Corresponding editor: Pierre-Olivier Antoine.

* Corresponding authors.

E-mail addresses: georgios.georgalis@pim.uzh.ch (G.L. Georgalis), massimo.delfino@unito.it (M. Delfino).

formally describe for the first time all the chelonian remains collected so far (a few specimens are exposed on the rock surface of the site) from the palaeoisland of Scontrone, which document the presence of at least two non-marine turtle lineages. The potential affinities and biogeographic origins of the new turtles from the Scontrone palaeoisland are discussed.

Geological settings

The fossiliferous locality of Scontrone is situated in the Parco Nazionale d'Abruzzo, Lazio e Molise, in Abruzzo region, Central Italy (41°45'15.54"N, 14°02'13.14"E; Fig. 1). The Scontrone fossil site consists of tidal-flat aeolian calcarenites at the base of the *Lithothamnion* Limestone, a Miocene carbonate ramp widespread in the central-southern Apennines (Savorelli et al., 2017). More details about the geology of the locality can be found in Patacca et al. (2008a), Patacca et al. (2013).

The locality is ubiquitously considered to pertain to a palaeoisland. The age of Scontrone is ascribed to the early Tortonian stage of the late Miocene, ca. 9 Ma (Patacca et al., 2013; Savorelli et al., 2017). Associated mammal fauna of Scontrone includes erinaceids, rodents, and hoplitomerycids (Mazza and Rustioni, 1996, 2008, 2011; Savorelli et al., 2007; Van Der Geer, 2014), while giant non-flying birds are also present (Pavia et al. 2017). Among reptiles, only a crocodylian (cf. *Crocodylus* sp.) has been described so far from Scontrone (Delfino and Rossi, 2013).



Fig. 1. The late Miocene (Tortonian) fossiliferous locality in Scontrone, Abruzzo, Italy. **A.** General panoramic view with the site (covered by a sliding protection) in the foreground. **B.** Close up with fossils still embedded in the rock.

Material and methods

All material described herein is permanently curated at the collections of the Centro di Documentazione Paleontologica “*Hoplitomeryx*”, Scontrone, Italy. Taxonomy and nomenclature of turtle clades follows Joyce et al. (2020), Joyce et al. (2021).

Institutional abbreviation: SCT, Centro di Documentazione Paleontologica “*Hoplitomeryx*”, Scontrone, Italy.

Systematic palaeontology

Testudines Batsch, 1788 [Joyce et al., 2020]
 Pan-Testudinoidae Joyce, Parham, and Gauthier, 2004 [Joyce et al., 2021]
 Testudinoidae Fitzinger, 1826 [Joyce et al., 2021]
 Geoemydidae Theobald, 1868 [Joyce et al., 2021]
 Genus *Mauremys* Gray, 1869
Mauremys sp.

Figs. 2, 3

Material: SCT 371a (a neural), SCT 371b (two costals and nine shell fragments); SCT 26a (a costal); SCT 26b (a costal); SCT 127a (a fragmentary costal); SCT 43 (five costal fragments and a bridge peripheral); SCT 256 (a peripheral); SCT 285 (a peripheral); SCT 258 (a peripheral); SCT 421 (a peripheral); SCT 416 (a peripheral); SCT 463 (a peripheral); SCT 218 (three peripherals); SCT 422 (a bridge peripheral); SCT 423 (fragment of a bridge peripheral); SCT 140a (a fragmentary ?suprapygale); SCT 233 (right epiplastron); SCT 214b (a left epiplastron); SCT 251 (a right epiplastron); SCT 334 (a right epiplastron fragment); SCT 253 (a fragmentary entoplastron); SCT 301 (a right hyoplastron); SCT 535 (a right hyoplastron on a slab); SCT 249 (a left hyoplastron); SCT 248 (a fragment of hyoplastron or hypoplastron); SCT 339 (a fragment of hyoplastron or hypoplastron); SCT 396 (a fragment of hyoplastron or hypoplastron); SCT 270 (a right hypoplastron); SCT 310 (a partial left hypoplastron); SCT 358 (a fragmentary hyo-hypoplastron); SCT 349 (a right xiphiplastron on a slab); SCT 239 (a left xiphiplastron); SCT 259 (several shell fragments on matrix); SCT 119 (an ilium).

Description

Carapace: The available carapace material consists of a neural, several costals and peripherals, and a probable suprapygale (Fig. 2). The single available neural is SCT 371a (Fig. 2(A)); it is not complete, as its anterolateral edge is broken off, but it is clear that the original general shape was hexagonal, with a concave anterior edge, and posterolateral edges much longer than the anterolateral ones. Most costals are fragmentary (Fig. 2(B, C)). Costal SCT 127a shows the possible trace of a long plastral buttress developed across a few centimetres along its ventral surface. Most shell remains available are peripherals, including bridge ones. SCT 421 is a peripheral pertaining to the beginning or the end of the bridge. Its external surface is characterized by a pleuro-marginal sulcus situated far from the costo-peripheral suture; it bears an intermarginal sulcus. SCT 416 is a complete peripheral originating from the anterior or posterior portion of the carapace, showing a clear intermarginal sulcus, but not a pleuro-marginal one. SCT 463 is a large-sized peripheral (probably peripheral I) that clearly shows the presence of a pleuro-marginal sulcus, as well as an intermarginal one (Fig. 2(F, G)). The peripherals of the bridge are relatively thin if compared with most of the plastral elements. SCT 218 consists of three peripherals of which the central one (which is also the most complete) seems to pertain to the left front of the bridge (Fig. 2(D, E)). It possesses pleuro-marginal and intermarginal furrows; prior to the beginning of the bridge cavity, at its




Fig. 2. Carapace elements of *Mauremys* sp. from the late Miocene (Tortonian) of Scontrone, Abruzzo, Italy. **A.** Neural SCT 371a in dorsal view. **B.** Costal SCT 26a in dorsal view. **C.** Costal SCT 26b in dorsal view. **D, E.** Three left bridge peripherals SCT 218 in anteroventral (D) and anterodorsal (E) views. **F, G.** Peripheral SCT 463 in dorsal (F) and ventral (G) views. Scale bars: 10 mm.

ventral surface, there is an elongated depression that could house the musk duct foramen (musk pore). SCT 140a is a fragmentary element that could represent a suprapygal. It is arched in cross-section, with sutures preserved on its ?posterior and ?left sides. A visible longitudinal sulcus is interpreted as the sulcus between

the last (likely XII) marginals, whereas a transversal sulcus is interpreted as the sulcus between the last marginals and the fifth vertebral scute.

Plastron: The plastral anatomy is relatively better known than that of the carapace, with plastron represented by epiplastra, an




Fig. 3. Plastron of *Mauremys* sp. from the late Miocene (Tortonian) of Scontrone, Abruzzo, Italy. **A–D.** Left epiplastron SCT 214b in dorsal (A), ventral (B), drawing of dorsal (C), and drawing of ventral (D) views. **E–I.** Right epiplastron SCT 233 in ventral (E), dorsal (F), anterior (G), drawing of ventral (H), and drawing of dorsal (I) views. **J, K.** Right hyoplastron SCT 535 on slab (J) and its drawing (K). **L, M.** Right xiphiplastron SCT 349 on slab, in lateral (L) and dorsal (M) views. **N, O.** Left xiphiplastron SCT 239 in ventral (N) and drawing of ventral (O) view. Scale bars: 10 mm.

entoplastron, hyo-hypoplastra, and xiphiplastron (Fig. 3). SCT 214b is a left epiplastron characterized by a slightly developed epiplastral lip (Fig. 3(A–D)). The latter is medially concave (well visible in anterior view). The gularo-humeral sulcus laterally corresponds

to a distinct notch on the epiplastron (corresponding also to the lateral edge of the epiplastral lip). The gularo-humeral sulcus intersects the epi-entoplastral suture ca. 7 mm from the inter-epiplastral suture indicating that the gulars significantly pene-

trated the entoplastron. There is no trace of a gular pocket, but rather only a weak depression posterior to the gular. SCT 251 is an incomplete right epiplastron, with an overall morphology congruent with that of SCT 214b, but slightly larger and thicker, being approximately 46 mm wide (due to erosion, this mediolateral width could be slightly larger in life). SCT 233 and SCT 334 correspond to right epiplastra (Fig. 3(E–I)). SCT 253 is a fragmentary entoplastron. Due to its fragmentary status, it is difficult to assess its exact shape, but it seems that it was wider than long. The sulci visible on the external surface clearly indicate that the gulars covered the anterior tip of the entoplastron and that the pectorals covered its posterior portion (i.e., the humero-pectoral sulcus crosses the entoplastron). On its internal surface there is a clearly elongated xiphoid scar. SCT 301 is a fragmentary right hyoplastron, partly covered by matrix (especially at the sutures with the epiplastron and entoplastron). Its anteroposterior length is ca. 67 mm long, whereas its maximum thickness reaches the 9.7 mm. This specimen preserves the humero-pectoral and pectoro-abdominal sulci. The position of the humero-pectoral sulcus clearly indicates that it crossed the entoplastron quite far from its posterior edge. The humero-pectoral sulcus is anteriorly concave and laterally terminates in correspondence at the weak, but still evident, notch in the lateral side of the anterior lobe of the plastron. The pectoro-abdominal sulcus is anteriorly convex and medially terminates far from the hyo-hyoplastral suture; its lateral termination cannot be evaluated due to preservational reasons. SCT 535 is a right hyoplastron on a slab (Fig. 3(J, K)). The humero-pectoral sulcus is clearly concave anteriorly. The pectoro-abdominal sulcus is equally undulating but is convex anteriorly in the central and middle portions of the hyoplastron, while it forms a deep anterior concavity close to its lateral edge. SCT 249 is a left hyoplastron whose external surface is so damaged that no sulci are visible. SCT 310 is a left hyoplastron that is only partially preserved. On its external surface, the abdomino-femoral forms an anterior convexity close to the lateral edge of the element. SCT 310 is slightly thickened (13.7 mm instead of 9.4 mm) at around its medial suture, in correspondence to the dorsal projection for the bridge buttress. Some further fragmentary elements are partly embedded in matrix but a definite identification cannot be made, such as the possible right hypoplastron SCT 270 and several fragments of hypoplastra or hypoplastra (SCT 248, SCT 339, and SCT 396) that show only their visceral surfaces. SCT 349 is a right xiphiplastron on a slab, situated along with an unidentified bone (Fig. 3(L, M)). This element shows only the visceral surface. Its lateral edge has a modest notch, corresponding to the femoro-anal sulcus, and a deep anal notch. The posterolateral edge of the xiphiplastron is rounded. The structure of the anterior edge of the element indicates the presence of a suture (not of a hinge). The thickness of the xiphiplastron, in the area close to its anterior suture, is 12 mm. SCT 239 is an incomplete left xiphiplastron, missing its anterior portion (Fig. 3(N, O)).

Appendicular skeleton: Only a single appendicular element is preserved, i.e., the ilium SCT 119. The glenoid cavity is not visible, due to preservational reasons, but the area surrounding it is well preserved. The distal (dorsal) edge is poorly preserved but it seems that some residues of concretion are hosted in a lateral, elongated concavity.

Remarks: The overall shape of the preserved skeletal elements along with the probable presence of musk duct foramina is consistent with a referral of the material to Geoemydidae (Claude and Tong, 2004; Garbin et al., 2018; Joyce et al., 2021). Furthermore, the paired marginals XII entering a probable suprapygal, the gulars reaching and overlapping the entoplastron, the humero-pectoral sulcus crossing the entoplastron, as well as

the deep anal notch of the xiphiplastron, suggest a referral to the genus *Mauremys* (Hervet, 2000; Claude et al., 2007).

Pan-Testudinidae Joyce, Parham, and Gauthier, 2004 [Joyce et al., 2021]

Testudinidae Gray, 1825 [Joyce et al., 2021]

Testudinidae indet. (large form)

Figs. 4–7

Material: SCT 118 (a partial posterior portion of a plastron partly preserved on a slab); SCT 127b–d (three plastral fragments); SCT 216a (a large shell fragment/unidentified laminar bone); SCT 237 (a ?pectoral girdle on a slab); SCT 163 (a pectoral girdle); SCT 203 (a fragment of a ?pubis); SCT 308 (a humerus); SCT 287 (a fragment of an ilium); SCT 236 (a fragment of a ?humerus); SCT 64 (a caudal vertebra).

Description

Plastron: No carapace material is present and therefore our knowledge on the shell anatomy of this taxon is based exclusively on its plastral remains (Figs. 4, 5). SCT 118 preserves the right hypoplastron and xiphiplastron (Fig. 4). Most of its external surface is covered by sediment but the exposed areas are rugose due to a prominent porosity (in the posterior tip of the xiphiplastron) or an irregular net of fine grooves (in the anterolateral sector of the hypoplastron). On the exposed area there is no evidence of sulci left by the scutes. The internal surface of both elements is relatively smooth. In the hypoplastron, its posterior edge and part of its posterolateral area are characterized by a long and nearly horizontal lateral buttress. The lateral edges of both the hypoplastron and xiphiplastron is not preserved and therefore, the original shape of the lateral edge of the posterior plastral lobe cannot be assessed. The xiphiplastron is nearly complete. This right xiphiplastron is rather thin (3 mm) along its sagittal suture with the (missing) left xiphiplastron but is comparatively much thicker at its posterior end, where a sort of “pad” nearly reaches 30 mm of thickness (the presence of fractures hinders a precise measurement of its thickness). Both the dorsal (visceral) and the ventral (external) surfaces of this posterior “pad” are distinctly convex, but it is not clear whether the ventral convexity reflects the original morphology or whether it is due to preservational reasons. However, this convexity, along with the general structure of the xiphiplastron and that of the hypoplastron, impart to the posterior lobe of the plastron a marked ventral concavity that possibly extended on the anterior lobe as well. The sagittal suture of this right xiphiplastron with its missing left counterpart is ca. 80 mm long; the suture with the hypoplastron is ca. 11 mm long, but it should have been slightly longer as both elements are laterally incomplete. Specimens SCT 127b–d seem also to represent plastral fragments (Fig. 5). None of them shows any sulcus on the external surface. In these plastral fragments, the external surface possesses irregular furrows (as in SCT 118), whereas the internal surface is rather smooth, with the exception of possible vascular sulci.

Appendicular skeleton: SCT 237 is a slab with a fragment of a possible pectoral girdle, preserving the glenoid cavity and the base of both the scapula and acromion (Fig. 6(A)). The rugose area close to the glenoid cavity could represent the contact surface with the coracoid. If so, the acromion is a slightly flattened structure that is a little thicker than the scapula. SCT 163 is a morphologically congruent element with SCT 237, but being isolated, it is more informative (Fig. 6(B)). Both the scapula and the acromion preserve only the area close to the glenoid cavity, which is filled by matrix. The contact surface with the coracoid is located on the ventral half of the acromion, close to the glenoid cavity. This surface is sur-




Fig. 4. Plastron of Testudinidae indet. (large form) from the late Miocene (Tortonian) of Scontrone, Abruzzo, Italy. Right hypoplastron and xiphiplastron SCT 118 in dorsal (A), ventral (B), and posterolateral (C) views. Scale bar: 50 mm.

mounted by a distinct tuberosity that is prolonged in a sort of ridge along the dorsal edge of the acromion. Both the scapula and the acromion are flattened and weakly concave on the surface (as the point where the contact surface is), but are slightly convex on the other side. A marked concavity corresponds to the above described tuberosity on the other side of the bone. SCT 203 is a large element that could represent the proximal ramus of a partial pubis (Fig. 6(C, D)). It possesses a triradiate structure with a broad and comparatively thick base distally, thinning into an arched front laterally, which continues into two processes that are broken off. One of the two sides of the thick base is rounded, the other is ridged.

SCT 308 is a fragmentary humerus preserving part of its distal portion, as the distal articulations are missing (Fig. 7(A, B)). The maximum preserved length of the specimen is of 43 mm. The ectepicondylar groove for the radial nerve is well visible. The sur-

face opposite to the groove has, towards the distal epiphysis, a longitudinal modest convexity delimited on both sides by two weak concavities. SCT 236 is a rather fragmentary specimen that could possibly also pertain to a humerus.

SCT 287 could probably represent the dorsal epiphysis of a very stout and large-sized right ilium (Fig. 7(C, D)). The epiphysis itself is not complete and the posterodorsal process is better preserved than the anterior one. The minimum diameters of the preserved portion are 28 and 17 mm, respectively.

Vertebra: SCT 64 is a partially preserved and slightly deformed caudal vertebra (centrum length = 26 mm) (Fig. 7(E, F)). The neural arch is severely damaged, though the elongate and slightly tilted prezygapophyseal facets are still visible. Part of the left postzygapophyseal facet is also present. The left transverse process in approximately horizontal (pointing ventrally only rather slightly) and laterally directed. It is flattened but dorsally provided by a



Fig. 5. Plastron of Testudinidae indet. (large form) from the late Miocene (Tortonian) of Scontrone, Abruzzo, Italy. Plastral fragment SCT 127b in dorsal (A) and ventral (B) views. Scale bar: 20 mm.




Fig. 6. Appendicular elements of Testudinidae indet. (large form) from the late Miocene of Scontrone, Abruzzo, Italy. **A.** ?Pectoral girdle SCT 237 on a slab. **B.** Pectoral girdle SCT 163. **C, D.** Partial ?pubis SCT 203 in dorsal (C) and ventral (D) views. Scale bars: 10 mm.

modest keel. The neural canal is proportionally small. The vertebra is procoelous and is characterized by two separated articular facets both anteriorly and posteriorly. The cotyle faces anteroventrally, whereas the condyle posterodorsally. The cotylar surfaces are separated by an evident notch. The ventral surface of the centrum is flattened, though a slight longitudinal keel is present at its midline.

Remarks: The presence of developed posterior buttresses on the plastron, along with the general shape of the elements denote a referral to testudinids (Vlachos, 2018). More particularly, within testudinids, the presence of such long and well-developed buttresses in the posterior portion of the plastron would hint for affinities with geocheilons. Within geocheilons, prominent posterior buttresses are characteristic of the




Fig. 7. Appendicular and vertebral elements of Testudinidae indet. (large form) from the late Miocene of Scontrone, Abruzzo, Italy. **A, B.** Partial humerus SCT 308 in posterior (A) and anterior (B) views. **C, D.** Partial right ilium SCT 287 in anterior (C) and posterior (D) views. **E, F.** Caudal vertebra SCT 64 in left lateral (E) and ventral (F) views. Scale bars: 10 mm.

European genus *Titanochelon* Pérez-García and Vlachos, 2014 (Pérez-García and Vlachos, 2014 (Pérez-García and Vlachos, 2014; Vlachos and Rabi, 2018)). However, more material is definitely needed in order to conclude whether the Scontrone tortoise has affinities with *Titanochelon* or even with some African form, such as *Centrochelys* Gray, 1872 (which also belongs to geocheilons). In any case, the size of the preserved elements denotes that they pertain to a relatively large form with a relatively short posterior lobe of the plastron, though any estimations about its probable carapace or plastron length would be speculative.

Testudines indet.

Material: SCT 127e–h (four shell fragments), SCT 5, SCT 214c, SCT 214d, SCT 289, SCT 298, SCT 302, SCT 360, SCT 378, SCT 472, SCT 474, SCT 408, SCT 485, SCT 497, SCT 498, SCT 500, SCT 501, SCT 503, SCT 508, SCT 509, SCT 511, SCT 512 (21 shell fragments), and SCT 106, SCT 387, and SCT 394 (three ?shell fragments).

Remarks. Besides the above described specimens, the assemblage comprises also several shell fragments that cannot be more precisely identified. Probably these could also pertain to

any of the two testudinoid forms described above, though the presence of an additional taxon cannot be ruled out with certainty. Nevertheless, affinities with trionychids, which are otherwise present in the Tortonian of Italy (Georgalis and Joyce, 2017; Georgalis et al., 2020), can be readily discarded, as the Scontrone shell fragments lack the characteristic sculpturing pattern of that turtle group.

Discussion

The Scontrone palaeoisland

Although not so well known as the nearby Gargano palaeoisland, Scontrone is characterized by an astonishing endemic fauna, comprising, among others, large erinaceids, the peculiar hoplitomerycids, and large non-flying birds (Van Der Geer, 2014; Pavia et al., 2017; Savorelli et al., 2017). Several of these insular taxa have congeneric relatives only in Gargano and nowhere else in the world, such as the iconic large erinaceid *Deinogalerix* Freudenthal, 1972 (Savorelli et al., 2017). Hoplitomerycids are also known exclusively from Gargano and Scontrone: the presence of different but still closely related genera in each palaeoisland, i.e., *Hoplitomeryx* Leinders, 1984, in Gargano, and *Scontromeryx* Van Der Geer, 2014, in Scontrone (Leinders, 1984; Mazza and Rustioni,

1996, 2011; Mazza, 2013; Van Der Geer, 2014) or of a single genus (*Hoplitomeryx*) in both palaeoislands is still debated (Mazza et al., 2016). The large non-flying anseriform bird *Garganornis ballmanni* Meijer, 2014, is also found exclusively in Gargano and Scontrone, notably being represented by conspecific finds in both palaeoislands (Meijer, 2014; Pavia et al., 2017). What appears rather important is the fact that, unlike Gargano that represents a timespan through part of the Neogene, the Scontrone palaeofauna is strictly confined to the early Tortonian, offering thus a definite stratigraphic accuracy and a unique glimpse into insular palaeoenvironments of the Central Mediterranean area of that time.

The origins and exact timing of dispersal of the Scontrone faunal assemblage have been a matter of debate, with different scenarios variously proposed (e.g., Abbazzi et al., 1996; Mazza and Rustioni, 1996, 2008; Masini et al., 2002, 2008, 2013; Rook et al., 2006; Freudenthal and Martín-Suárez, 2010). In general, the whole Abruzzo-Apulia Platform, to which Scontrone belonged, was isolated from Continental Europe at least since the early Oligocene and up to the latest Miocene (Mazza, 2013). In fact, it has been suggested that the last interconnection of the Platform was through a terrestrial land corridor in the Adriatic with northwestern Balkans during the early Oligocene, ca. 29 Ma (Patacca et al., 2008b; Mazza, 2013). It is thus not fully settled whether Gargano and Scontrone were colonized one or multiple times independently, though recent studies have provided strong evidence for the latter case, i.e., multiphasic colonization events (Rook et al., 2006; Savorelli and Masini, 2016; Savorelli et al., 2017). In particular, different scenarios have been proposed for the most famous inhabitants of the Abruzzo-Apulia Platform: for *Deinogalerix* it has been suggested that its origins should be traced in an early to middle Miocene ancestor (Savorelli et al., 2017), whereas for the peculiar hoplitomerycids that they arrived in the area already by the Oligocene (Mazza and Rustioni, 2011).

Palaeobiogeography of the Scontrone turtles

Taking into consideration the complex biogeographic history of the insular Abruzzo-Apulia faunas discussed above, it seems a rather plausible scenario to adhere that the geoemydid and testudinid ancestors of the Scontrone chelonians described herein are of Balkan or even Western Asian origin that arrived in the area during the Oligocene. For *Mauremys*, the possibility of an Oligocene dispersal from the Balkans to Scontrone palaeoisland is hindered by the rather poor and limited fossil record of this genus in the Paleogene of Europe (Hervet, 2004), and particularly in the Balkan Peninsula. In fact, the only Oligocene taxon from the Balkans (but also from the whole Eastern Europe) that could potentially pertain to *Mauremys* is *Mauremys strandi* (Szalai, 1934) from the early Oligocene of Romania (Szalai, 1934; Młynarski and Mészáros, 1963), however, the generic placement of this species is still far from clear. Also, two further Balkan taxa are *Palaeochelys rueckerti* Schleich, 1994, and *Palaeochelys turcica* Schleich, 1994, both from the late Oligocene or early Miocene of the European part of Turkey (Schleich, 1994), however, these are represented by rather incomplete material and their affinities with *Mauremys* are not yet resolved.

On the other hand, African origins have been envisaged for the sole other described reptilian inhabitant of Scontrone, i.e., the crocodylid cf. *Crocodylus*, which has been suggested to represent the product of a direct or indirect marine dispersal from Africa that colonized the palaeoisland of Scontrone during the Tortonian (Delfino and Rossi, 2013).

In any case, fossil record, molecular data, as well as direct observations of extant taxa, attest that *Mauremys* and, especially, large testudinids are well capable of dispersing throughout large marine distances (Gerlach et al., 2006; Vamberger et al., 2014; Kehlmaier

et al., 2019; Georgalis et al., 2020). As such, a direct terrestrial corridor would not be strictly necessary for these turtle taxa to reach Scontrone. Therefore, the new geoemydid and large testudinid from Scontrone could represent the product(s) of some dispersal(s) during the late Paleogene and/or Neogene, though we have to leave open the possibility whether such dispersal(s) took place from either Africa, or the Balkans or Central Europe. Nevertheless, in particular for the geoemydid, a Paleogene or early Neogene African origin seems unlikely, as this lineage apparently dispersed to Africa from Europe only during the Tortonian (Georgalis et al., 2020) and is totally absent from pre-Tortonian sediments of Afro-Arabia (Georgalis et al., 2021); therefore, if *Mauremys* reached Scontrone following a southern to northern route, then this would not predate the Tortonian, implying that geoemydids were only new-comers on the palaeoisland, similarly to the case of their crocodylian co-inhabitant (Delfino and Rossi, 2013).

The identification of *Mauremys* in the Scontrone assemblage adds to the known distribution of this genus in Italy, which consists of a ubiquitous faunal element throughout large part of the Neogene and Quaternary, being nevertheless, currently extinct from the area (Sacco, 1889; Portis, 1890, 1896; Kotsakis, 1981; Delfino and Bailon, 2000; Girotti et al., 2003; Chesi et al., 2007, 2009a; Colombero et al., 2017; Collareta et al., 2020; Georgalis et al., 2020). It further attests for a wider distribution of *Mauremys* throughout the Mediterranean during the Tortonian (Georgalis et al., 2016, 2020). As the present material cannot afford an identification to the species level, it cannot be determined whether the Scontrone geoemydid represents an endemic insular form or whether it is conspecific with any of the mainland European taxa, including the Italian extinct species, i.e., the late Miocene *Mauremys campanii* (Ristori, 1891), the Pliocene *Mauremys portisii* (Sacco, 1889), and the Plio-Pleistocene *Mauremys etrusca* (Portis, 1890).

As for the large-sized tortoise from Scontrone, its exact affinities are also hindered by its poor preservational status, which mirrors the one of the few, likewise large, testudinid remains coming from Gargano (Delfino, 2002). Large to giant tortoises were present in Europe throughout most of the Cenozoic (Lapparent de Broin, 2002; Georgalis and Kear, 2013; Pérez-García and Vlachos, 2014; Vlachos et al., 2020). Similarly, large-sized testudinids were also present during the Cenozoic of northern Africa, though their fossil record there is significantly more poorly documented in comparison with their European counterpart (Lapparent de Broin, 2002; Georgalis et al., 2021). In the Mediterranean Islands, testudinids have a relatively rich fossil record, that comprises an array of endemic forms during the late Neogene and Quaternary of several islands (e.g., Malta, Balearic Islands, Sardinia, Sicily), but all of these are from Plio-Pleistocene localities, thus postdating the Tortonian age of the Scontrone material (Leith Adams, 1877; Bate, 1914; Delfino, 2002; Abbazzi et al., 2008; Luján et al., 2017). This diversity and abundance of Mediterranean insular testudinids, strongly evokes the situation in different oceanic islands across the globe. There, several extant and recently (or at least relatively recently) extinct endemic testudinids flourished. These include the famous and diverse radiations in the Galapagos Islands (*Chelonoidis* Fitzinger, 1836; extant), the Mascarenes (*Cylindraspis* Fitzinger, 1836; extinct only few centuries ago), the Seychelles Archipelago and Madagascar (*Aldabrachelys* Loveridge and Williams, 1957; Quaternary, including also an extant taxon), the Caribbean Sea and the Bahamas Archipelago (*Chelonoidis*; Quaternary), and certain more poorly known taxa from the Canary Islands (“*Centrochelys*”; Plio-Pleistocene), the Cape Verde Islands (“*Centrochelys*”; Pleistocene), Indonesia (?*Megalochelys* Falconer and Cautley, 1837; Pleistocene), and Philippines (?*Megalochelys*; Pleistocene) (Günther, 1877; Van Denburgh, 1914; Arnold, 1979; Gerlach, 2004; Setiyabudi, 2009; Rhodin et al., 2015; Vlachos,

2018; Kehlmaier et al., 2019; Georgalis et al., 2021). A principal common feature of almost all of these endemic insular testudinid forms is the acquisition of a large size. These being said, the Scontrone testudinid could be a descendant of any of the late Paleogene or Neogene continental large forms that inhabited Europe and Afro-Arabia, or even of some smaller continental form that eventually evolved a large size during its insular isolation. More remains are definitely needed in order to decipher the exact taxonomic affinities and biogeographic origins of the Scontrone turtles.

We are still only at the beginning of comprehending these unique insular herpetofaunas on these now long-lost islands of the Central Mediterranean. Even though poorly preserved, the Gargano and Scontrone herpetofaunas hint towards interesting dispersal events and unique radiations. It is further evident that Central and Southern Italy played a dynamic role in the biogeography of turtles during the Tortonian, as is also exemplified by recently described coeval finds from Calabria, which could point towards the dispersal of geoemydids and trionychids from Europe to Africa through this part of the Central Mediterranean (Georgalis et al., 2020).

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We are grateful to Adelaide Rossi and the Soprintendenza Archeologia Belle Arti e Paesaggio dell'Abruzzo (Chieti, Italy) for having organized the retrieval of the fossil remains from Scontrone and promoting their study. MD thanks Paul P.A. Mazza who introduced him to the faunal assemblage of Scontrone and for having invited him for the field season 2002. GLG acknowledges funding by Forschungskredit of the University of Zurich, grant n° FK-20-110 and a former postdoctoral grant from the University of Torino. MD acknowledges funding by Fondi di Ateneo dell'Università di Torino (2019–2020), Generalitat de Catalunya (CERCA Program), and Spanish Agencia Estatal de Investigación (CGL2016-76431-P, AEI/FEDER, EU). The quality of the paper was improved by useful comments provided by Paul P.A. Mazza, the editor Gilles Escarguel, the associate editor Pierre-Olivier Antoine, and the reviewer Julien Claude.

References

- Abbazzi, L., Benvenuti, M., Boschian, G., Dominici, S., Masini, F., Mezzabotta, C., Piccini, L., Rook, L., Valleri, G., Torre, D., 1996. Revision of the Neogene and Pleistocene of the Gargano region (Apulia, Italy). The marine and continental succession and the mammal faunal assemblages in an area between Apricena and Poggio Imperiale (Foggia). *Memorie della Società Geologica Italiana* 51, 383–402.
- Abbazzi, L., Carboni, S., Delfino, M., Gallai, G., Lecca, L., Rook, L., 2008. Fossil vertebrates (Mammalia and Reptilia) from Capo Mannu Formation (Late Pliocene, Sardinia, Italy), with description of a new *Testudo* (Chelonii, Testudinidae) species. *Rivista Italiana di Paleontologia e Stratigrafia* 114, 19–132.
- Arnold, E.N., 1979. Indian Ocean giant tortoises: their systematics and island adaptations. *Philosophical Transactions of the Royal Society of London B* 286, 127–145.
- Ballmann, P., 1976. Fossile Vögel aus dem Neogen der Halbinsel Gargano (Italien), zweiter Teil. *Scripta Geologica* 38, 1–59.
- Bate, D.M.A., 1914. On remains of a gigantic land tortoise (*Testudo gymnesicus*, n. sp.) from the Pleistocene of Menorca. *Geological Magazine* 1, 100–107.
- Batsch, A.J.G.K., 1788. Versuch einer Anleitung, zur Kenntniß und Geschichte der Thiere und Mineralien, für akademische Vorlesungen entworfen, und mit den nöthigsten Abbildungen versehen, Erster Theil. Allgemeine Geschichte der Natur; besondere der Säugethiere, Vögel, Amphibien und Fische. Akademische Buchhandlung, Jena.
- Blain, H.-A., Fagoaga, A., Ruiz-Sánchez, F.J., Bisbal-Chinesta, J.F., Delfino, M., 2019. Latest Villafranchian climate and landscape reconstructions at Pirro Nord (southern Italy). *Geology* 47, 829–832.
- Chesi, F., Delfino, M., Abbazzi, L., Carboni, S., Lecca, L., Rook, L., 2007. New fossil vertebrate remains from San Giovanni di Sinis (Late Pleistocene, Sardinia): the last *Mauremys* (Reptilia, Testudines) in the central Mediterranean. *Rivista Italiana di Paleontologia e Stratigrafia* 113, 287–297.
- Chesi, F., Delfino, M., Rook, L., 2009a. Late Miocene *Mauremys* (Testudines, Geoemydidae) from Tuscany (Italy): evidence of terrapin persistence after a mammal turnover. *Journal of Paleontology* 83, 379–388.
- Chesi, F., 2009b. Il registro fossile italiano dei chelonii. Ph.D. Thesis, Università di Firenze, Firenze, Italy. 178 pp.
- Claude, J., Tong, H., 2004. Early testudinoid turtles from Saint-Papoul, France, with comments on the early evolution of modern Testudinoidea. *Oryctos* 5, 3–45.
- Claude, J., Tong, H., Suteethorn, V., 2007. Turtles from the late Eocene–early Oligocene of the Krabi Basin (Thailand). *Bulletin de la société géologique de France Série 1* (178), 305–316.
- Collareta, A., Casati, S., Zuffi, M.A.L., di Cencio, A., 2020. First authentic record of the freshwater turtle *Mauremys* from the Upper Pliocene of Italy, with a new occurrence of the rarely reported ichnotaxon *Thatchtelithichnus holmani*. *Carnets de Géologie* 20, 301–313.
- Colombero, S., Alba, D.M., D'Amico, C., Delfino, M., Esu, D., Giuntelli, P., Harzhauser, M., Mazza, P.P.A., Mosca, M., Neubauer, T.A., Pavia, G., Pavia, M., Villa, A., Carnevale, G., 2017. Late Messinian mollusks and vertebrates from Moncucco Torinese, north-western Italy. Paleoeological and paleoclimatological implications. *Palaeontologia Electronica* 20.1.10A, 1–66.
- Delfino, M., 2002. Erpetofaune italiane del Neogene e del Quaternario. Ph.D. Thesis, University of Modena and Reggio Emilia, Modena and Reggio Emilia (unpubl.).
- Delfino, M., Böhme, M., Rook, L., 2007. First European evidence for transcontinental dispersal of *Crocodylus* (late Neogene of southern Italy). *Zoological Journal of the Linnean Society* 149, 293–307.
- Delfino, M., Bailon, S., 2000. Early Pleistocene herpetofauna from Cava Dell' Erba and Cava Pirro (Apulia, Southern Italy). *Herpetological Journal* 10, 95–110.
- Delfino, M., Rossi, M.A., 2013. Fossil crocodylid remains from Scontrone (Tortonian, Southern Italy) and the late Neogene Mediterranean biogeography of crocodylians. *Geobios* 46, 25–31.
- Falconer, H., Cautley, P.T., 1837. On additional fossil species of the order Quadrumana from the Siwalik Hills. *Journal of the Asiatic Society of Bengal* 6, 354–360.
- Fitzinger, L.J., 1826. Neue Classification der Reptilien, nach ihren Natürlichen Verwandtschaften nebst einer Verwandtschafts-Tafel und einem Verzeichnisse der Reptilien-Sammlung des k.k. Zoologischen Museum zu Wien. J.G. Hübner Verlage, Wien.
- Fitzinger, L.J., 1836. Entwurf einer systematischen Anordnung der Schildkröten nach den Grundsätzen der natürlichen Methode. *Annalen des Wiener Museums der Naturgeschichte* 1, 104–128.
- Freudenthal, M., 1971. Neogene vertebrates from the Gargano Peninsula, Italy. *Scripta Geologica* 3, 1–10.
- Freudenthal, M., 1972. *Deinogalerix koenigswaldi* nov. gen., nov. sp., a giant insectivore from the Neogene of Italy. *Scripta Geologica* 14, 1–19.
- Freudenthal, M., 1976. Rodent stratigraphy of some Miocene fissure fillings in Gargano (prov. Foggia, Italy). *Scripta Geologica* 27, 1–23.
- Freudenthal, M., 1985. Cricetidae (Rodentia) from the Neogene of Gargano (prov. Foggia, Italy). *Scripta Geologica* 77, 29–75.
- Freudenthal, M., Martín-Suárez, E., 2010. The age of immigration of the vertebrate faunas found at Gargano (Apulia, Italy) and Scontrone (l'Aquila, Italy). *Comptes Rendus Palevol* 9, 95–100.
- Freudenthal, M., Van den Hoek Ostende, L.W., Martín-Suárez, E., 2013. When and how did the *Mikrotia* fauna reach Gargano (Apulia, Italy)? *Geobios* 46, 105–109.
- Garbin, R.C., Ascarrunz, E., Joyce, W.G., 2018. Polymorphic characters in the reconstruction of the phylogeny of geoemydid turtles. *Zoological Journal of the Linnean Society* 184, 896–918.
- Georgalis, G.L., Arca, M., Rook, L., Tuveri, C., Delfino, M., 2019. A new colubroid snake (Serpentes) from the early Pleistocene of Sardinia, Italy. *Bollettino della Società Paleontologica Italiana* 58, 277–294.
- Georgalis, G.L., Insacco, G., Rook, L., Spadola, F., Delfino, M., 2020. Turtle remains from the late Miocene of the Cessaniti area, southern Italy – insights for a probable Tortonian chelonian dispersal from Europe to Africa. *Swiss Journal of Paleontology* 139, 1.
- Georgalis, G.L., Joyce, W.G., 2017. A review of the fossil record of Old World turtles of the clade *Pan-Trionychidae*. *Bulletin of the Peabody Museum of Natural History* 58, 115–208.
- Georgalis, G.L., Kear, B.P., 2013. The fossil turtles of Greece: An overview of taxonomy and distribution. *Geobios* 46, 299–311.
- Georgalis, G.L., Macaluso, L., Delfino, M., 2021. A review of the fossil record of Afro-Arabian turtles of the clade *Testudinoidea*. *Bulletin of the Peabody Museum of Natural History* 62, 43–78.
- Georgalis, G.L., Villa, A., Vlachos, E., Delfino, M., 2016. Fossil amphibians and reptiles from Plakias, Crete: a glimpse into the earliest late Miocene herpetofaunas of southeastern Europe. *Geobios* 49, 433–444.
- Gerlach, J., 2004. Giant tortoises of the Indian Ocean: the genus *Dipsochelys* inhabiting the Seychelles Islands and the extinct giants of Madagascar and the Mascarenes. *Frankfurter Beiträge zur Naturkunde* 21. Edition Chimaira, Frankfurt am Main.
- Gerlach, J., Muir, C., Richmond, M.D., 2006. The first substantiated case of trans-oceanic tortoise dispersal. *Journal of Natural History* 40, 2403–2408.

- Girrotti, O., Capasso Barbato, L., Esu, D., Gliozzi, E., Kotsakis, T., Martinetto, E., Petronio, C., Sardella, R., Squazzini, E., 2003. The section of Torre di Picchio (Terni, Umbria, Central Italy): a Villafranchian site rich in vertebrates, molluscs, ostracods and plants. *Rivista Italiana di Paleontologia e Stratigrafia* 109, 77–98.
- Gray, J.E., 1825. A synopsis of the genera of reptiles and amphibia, with a description of some new species. *Annals of Philosophy* 10, 193–217.
- Gray, J.E., 1869. Description of *Mauremys lanjaria*, a new freshwater tortoise. In: *Proceedings of the Zoological Society of London 1869*, pp. 499–500.
- Gray, J.E., 1872. Appendix to the catalogue of shield reptiles in the collection of the British Museum. Part I. Testudinata (tortoises). London: British Museum, 28 p.
- Günther, A., 1877. The gigantic land-tortoises (living and extinct) in the collection of the British Museum. Taylor and Francis, London.
- Herridge, V.L., Lister, A.M., 2012. Extreme insular dwarfism evolved in a mammoth. *Proceedings of the Royal Society B: Biological Sciences* 279, 3193.
- Hervet, S., 2000. Tortues du Quaternaire de France : critères de détermination, répartition chronologique et géographique. *Mésogée* 58, 3–47.
- Hervet, S., 2004. Systématique du groupe « *Palaeochelys* sensu lato–*Mauremys* » (Chelonii, Testudinoidea) du Tertiaire d'Europe occidentale : principaux résultats. *Annales de Paléontologie* 90, 13–78.
- Joyce, W.G., Anquetin, J., Cadena, E.-A., Claude, J., Danilov, I.G., Evers, S.W., Ferreira, G.S., Gentry, A.D., Georgalis, G.L., Lyson, T.R., Pérez-García, A., Rabi, M., Sterli, J., Vitek, N.S., Parham, J.F., 2021. A nomenclature for fossil and living turtles using phylogenetically defined clade names. *Swiss Journal of Palaeontology* 140, 5.
- Joyce, W.G., Parham, J.F., Anquetin, J., Claude, J., Danilov, I.G., Iverson, J.B., Kear, B., Lyson, T.R., Rabi, M., Sterli, J., 2020. Testudines. In: de Queiroz, K., Cantino, P.D., Gauthier, J.A. (Eds.), *Phylogeny – A Companion to the PhyloCode*. CRC Press, Boca Raton, pp. 1049–1051.
- Joyce, W.G., Parham, J.F., Gauthier, J.A., 2004. Developing a protocol for the conversion of rank-based taxon names to phylogenetically defined clade names, as exemplified by turtles. *Journal of Paleontology* 78, 989–1013.
- Kehlmaier, C., Graciá, E., Campbell, P.D., Hofmeyr, M.D., Schweiger, S., Martínez-Silvestre, A., Joyce, W., Fritz, U., 2019. Ancient mitogenomics clarifies radiation of extinct Mascarene giant tortoises (*Cylindraspis* spp.). *Scientific Reports* 9, 17487.
- Kotsakis, T., 1981. Gli anfibi e i rettili dei Pleistocene del Lazio (Italia Centrale). *Geologica Romana* 20, 57–67.
- Lanza, B., 1984. Sul significato biogeografico delle isole fossili, con particolare riferimento all'arcipelago pliocenico della Toscana. *Atti della Società italiana di scienze naturali e del Museo civico di storia naturale di Milano* 125, 145–158.
- Lapparent de Broin, F. de, 2002. A giant tortoise from the Late Pliocene of Lesbos Island (Greece) and its possible relationships. *Annales Géologiques des Pays Helléniques* 39, 99–130.
- Leinders, J., 1984. *Hoplitomerycidae* fam. nov. (Ruminantia, Mammalia) from Neogene fissure fillings in Gargano (Italy). *Scripta Geologica* 70, 1–68.
- Leith Adams, A., 1877. On gigantic land-tortoises and a small freshwater species from the ossiferous caverns of Malta, together with a list of their fossil fauna; and a note on chelonian remains from the rock cavities of Gibraltar. *Quarterly Journal of the Geological Society of London* 33, 177–191.
- Loveridge, A., Williams, E., 1957. Revision of the African tortoises and turtles of the suborder Cryptodira. *Bulletin of the Museum of Comparative Zoology* 115, 163–557.
- Luján, A.H., Alcover, J.A., Ivanov, M., Torres, E., Alba, D.M., 2017. Revisión taxonómica de “*Testudo*” *gymnesica* Bate, 1914 (Testudines, Testudinidae) a partir de la descripción del material tipo de Menorca (Islas Baleares). *Spanish Journal of Palaeontology* 32, 261–278.
- Masini, F., Bonfiglio, L., Petruso, D., Marra, A.C., Abbazzi, L., Delfino, M., Fanfani, F., Torre, D., 2002. The role of coastal areas in the Neogene-Quaternary mammal island populations of central Mediterranean. *Biogeographia* 23, 165–189.
- Masini, F., Fanfani, F., 2013. *Apulogalerix pusillus* nov. gen., nov. sp., the small-sized Galericiinae (Erinaceidae, Mammalia) from the “Terre Rosse” fissure filling of the Gargano (Foggia, South-Eastern Italy). *Geobios* 46, 89–104.
- Masini, F., Petruso, D., Bonfiglio, L., Mangano, G., 2008. Origination and extinction patterns of mammals in three central Western Mediterranean islands from the Late Miocene to Quaternary. *Quaternary International* 182, 63–79.
- Masini, F., Rinaldi, P.M., Petruso, D., Surdi, G., 2010. The Gargano Terre Rosse faunas: an overview. *Rivista Italiana di Paleontologia e Stratigrafia* 116, 421–435.
- Masini, F., Rinaldi, P., Savorelli, A., Pavia, M., 2013. A new small mammal assemblage from the M013 Terre Rosse fissure filling (Gargano, South-Eastern Italy). *Geobios* 46, 49–61.
- Mazza, P.P.A., 2013. *Hoplitomerycidae* (Ruminantia; Late Miocene, Central-Southeastern Italy): Whom and where from? *Geobios* 46, 511–520.
- Mazza, P., Rossi, M.A., Rustioni, M., Agostini, S., Masini, F., Savorelli, A., 2016. Observations on the postcranial anatomy of *Hoplitomeryx* (Mammalia, Ruminantia, Hoplitomerycidae) from the Miocene of the Apulia Platform (Italy). *Palaeontographica Abteilung A* 307, 105–147.
- Mazza, P., Rustioni, M., 1996. The Turolian artiodactyls from Scontrone (Abruzzi, central Italy). *Bollettino Società Paleontologia Italiana* 35, 93–106.
- Mazza, P., Rustioni, M., 2008. Processes of island colonization by Oligo-Miocene land mammals in the central Mediterranean: New data from Scontrone (Abruzzo, Central Italy) and Gargano (Apulia, Southern Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology* 267, 208–215.
- Mazza, P., Rustioni, M., 2011. Five new species of *Hoplitomeryx* from the Neogene of Abruzzo and Apulia (central and southern Italy) with revision of the genus and of *Hoplitomeryx matthei* Leinders, 1983. *Zoological Journal of the Linnean Society* 163, 1304–1333.
- Meijer, H.J.M., 2014. A peculiar anseriform (Aves: Anseriformes) from the Miocene of Gargano (Italy). *Comptes Rendus Palevol* 13, 19–26.
- Młynarski, M., Mészáros, M., 1963. Systematic position of *Clemmys strandi* (Szalai, 1934) (Testudines, Emydidae), from the Upper Oligocene of Cluj (Romania). *Acta Zoologica Cracoviensia* 8, 327–334.
- Palombo, M.R., Rozzi, R., Bover, P., 2013. The endemic bovids from Sardinia and the Balearic Islands: state of the art. *Geobios* 46, 127–142.
- Patacca, E., Scandone, P., Carnevale, G., 2013. The Miocene vertebrate-bearing deposits of Scontrone (Abruzzo, Central Italy): Stratigraphic and paleoenvironmental analysis. *Geobios* 46, 5–23.
- Patacca, E., Scandone, P., Mazza, P., 2008a. The Miocene land-vertebrate fossil site of Scontrone (Central Apennines). *Bollettino della Società Geologica Italiana* 127, 51–73.
- Patacca, E., Scandone, P., Mazza, P., 2008b. Oligocene migration path for Apulia macromammals: the Central-Adriatic bridge. *Bollettino della Società Geologica Italiana* 127, 337–355.
- Pavia, M., 2013. The Anatidae and Scolopacidae (Aves: Anseriformes, Charadriiformes) from the late Neogene of Gargano, Italy. *Geobios* 46, 43–48.
- Pavia, M., Meijer, H.J.M., Rossi, M.A., Göhlich, U.B., 2017. The extreme insular adaptation of *Garganornis ballmanni* Meijer, 2014: a giant Anseriformes of the Neogene of the Mediterranean Basin. *Royal Society Open Science* 4, 160722.
- Pérez-García, A., Vlachos, E., 2014. New generic proposal for the European Neogene large testudinids (Cryptodira) and the first phylogenetic hypothesis for the medium and large representatives of the European Cenozoic record. *Zoological Journal of the Linnean Society* 172, 653–719.
- Portis, A., 1890. I rettili pliocenici del Valdarno superiore e di alcune altre località plioceniche di Toscana. Le Monnier, Florence.
- Portis, A., 1896. Contribuzioni alla storia fisica del bacino di Roma e studii sopra l'estensione da darsi al Pliocene superiore. L. Roux, Rome.
- Rhodin, G.J.A., Thomson, S., Georgalis, G.L., Karl, H.-V., Danilov, I.G., Takahashi, A., De La Fuente, M.S., Bourque, J.S., Delfino, M., Bour, R., Iverson, J.B., Shaffer, H.B., Van Dijk, P.P., 2015. Turtles and tortoises of the world during the rise and global spread of humanity: first checklist and review of extinct Pleistocene and Holocene chelonians. *Chelonian Research Monographs* 5, 1–66.
- Ristori, G., 1891. I cheloniani fossili di Montebamboli e Casteani (Maremma, Toscana). *Atti della Società Toscana di Scienze Naturali, Processi Verbal* 7, 304–307.
- Rook, L., Abbazzi, L., Chesi, F., Delfino, M., Ferretti, M.P., Gallai, G., 2008. The Italian record of latest Miocene continental vertebrates. *Bollettino della Società Paleontologia Italiana* 47, 191–194.
- Rook, L., Gallai, G., Torre, D., 2006. Lands and endemic mammals in the Late Miocene of Italy: Constraints for paleogeographic outlines of Tyrrhenian area. *Palaeogeography, Palaeoclimatology, Palaeoecology* 238, 263–269.
- Rustioni, M., Mazza, P., Azzaroli, A., Boscagli, G., Cozzini, F., Di Vito, E., Masseti, M., Pisanò, A., 1992. Miocene vertebrate remains from Scontrone, National Park of Abruzzi, Central Italy. *Rendiconti Lincei. Scienze Fisiche e Naturali*, series 9 3, 227–237.
- Sacco, F., 1889. I Cheloni astiani del Piemonte. *Memorie della Reale Accademia delle Scienze di Torino* 39, 427–461.
- Savorelli, A., Masini, F., 2016. *Mystemys giganteus* nov. gen. nov. sp.: an enigmatic and rare critetid from the Terre Rosse M013 fissure filling (Gargano, Southeastern Italy). *Palaeontographica Abteilung A* 306, 1–23.
- Savorelli, A., Masini, F., Mazza, P.P.A., Rossi, M.A., Agostini, S., 2017. New species of *Deinogalerix* (Mammalia, Eulipotyphla) from the late Miocene of Scontrone (Abruzzo, central Italy). *Palaeontologia Electronica* 20.1.16A, 1–26.
- Setiyabudi, E., 2009. An early Pleistocene giant tortoise (Reptilia; Testudines; Testudinidae) from the Bumiayu area, Central Java, Indonesia. *Journal of Fossil Research* 42, 1–11.
- Szalai, T., 1934. Die fossilen Schildkröten Ungarns. *Folia Zoologica et Hydrobiologica* 6, 97–142.
- Theobald, W., 1868. Catalogue of reptiles in the museum of the Asiatic Society of Bengal. *Journal of the Asiatic Society, Extra Number* 1868, 8–88.
- Vamberger, M., Stuckas, H., Ayaz, D., Lymberakis, P., Široký, P., Fritz, U., 2014. Massive transoceanic gene flow in a freshwater turtle (Testudines: Geoemydidae: *Mauremys rivulata*). *Zoologica Scripta* 43, 313–322.
- Van Denburgh, J., 1914. Expedition of the California Academy of Sciences to the Galapagos Islands, 1905–1906. X. The gigantic land tortoises of the Galapagos Archipelago. *Proceedings of the California Academy of Sciences* 2, 203–374.
- Van der Geer, A., Lyras, G., de Vos, J., Dermitzakis, M., 2010. Evolution of Island Mammals: Adaptation and Extinction of Placental Mammals on Islands. Wiley-Blackwell, Oxford.
- Van Der Geer, A., 2014. Systematic revision of the family Hoplitomerycidae Leinders, 1984 (Artiodactyla: Cervioidea), with the description of a new genus and four new species. *Zootaxa* 3847, 1–32.
- Villier, B., Carnevale, G., 2013. A new skeleton of the giant hedgehog *Deinogalerix* from the Miocene of Gargano, southern Italy. *Journal of Vertebrate Paleontology* 33, 902–923.
- Vlachos, E., 2018. A review of the fossil record of North American turtles of the clade *Pan-Testudinoidea*. *Bulletin of the Peabody Museum of Natural History* 59, 3–94.
- Vlachos, E., Georgalis, G.L., Roussiakis, S., Böhme, M., Theodorou, G., 2020. The Pliocene tortoises (Testudines, Testudinidae) from the late Miocene of the South Balkans. *Journal of Vertebrate Paleontology* 39, e1711520.
- Vlachos, E., Rabi, M., 2018. Total evidence analysis and body size evolution of extant and extinct tortoises (Testudines: Cryptodira: Pan-Testudinidae). *Cladistics* 34, 652–683.